PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals

CAUTION: Keep away from children, domestic animals and pets. May be harmful if swallowed because it may reduce the clotting ability of blood and cause bleeding. Causes moderate eye irritation. Do not get in eyes, on skin or on clothing. Wear gloves when handling bait. Wash thoroughly with soap and water after applying and before eating, drinking, chewing gum or using tobacco. Remove and wash contaminated clothing before reuse.

FIRST AID

Have this label with you when obtaining treatment advice. If swallowed: Call a poison control center or doctor immediately for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin with plenty of cool water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Note to Physician or Veterinarian: Contains bromadiolone, an anticoagulant. For humans or pets that have ingested this product, or have obvious poisoning symptoms (bleeding) or prolonged prothrombin times, give Vitamin $\rm K_1$ by intramuscular or oral administration. Check prothrombin time every 3 days until values return to normal.

Environmental Hazards: This product is toxic to fish and wildlife. Predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten the bait. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Storage: Store in original container in a cool, dry place inaccessible to children and pets.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Container Disposal: Dispose of empty container in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



PARAFFIN BLOCKS

KILLS NORWAY RATS, ROOF RATS, HOUSE MICE AND WARFARIN-RESISTANT NORWAY RATS

NORWAY RATS & HOUSE MICE CAN CONSUME A LETHAL DOSE IN ONE NIGHT'S FEEDING WITH FIRST DEAD RODENTS APPEARING FOUR OR FIVE DAYS AFTER FEEDING BEGINS

 Active Ingredient: bromadiolone
 0.005%

 Inert Ingredients
 99.995%

 Total
 100.000%

KEEP OUT OF REACH OF CHILDREN

CAUTION: See side panel for additional precautionary statements.



Liphatech, Inc. 3600 W. Elm Street Milwaukee, WI 53209 (414) 351-1476

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL and follow all use directions and precautions. For information on this product (including health concerns, medical emergencies or pesticide incidents) call the National Pesticide Information Center at 1-800-858-7378.

IMPORTANT: Do not expose children, pets or other nontarget animals to rodenticides. To help prevent accidents:

- 1. Store product not in use in a location out of reach of children and pets.
- 2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
- 3. Dispose of product container, unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may be used to control Norway rats, roof rats and house mice. Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment. Do not broadcast bait.

Urban areas: This product may be used in and around homes, food processing facilities, industrial, commercial and public buildings; in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. This product may also be used in alleys and sewers.

Non-urban areas: This product may be used inside of homes and agricultural buildings.

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely find and consume bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors or walls, or in locations where rodents or their signs have been seen. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

Each block is scored to be broken into 2-oz. pieces.

Mice: Break block and apply one 2-oz. piece per placement. Space placements 8 to 12 feet apart. Maintain a constant supply of fresh bait for 15 days or until fresh signs of mouse activity cease.

Rats: Apply two to eight 2-oz. pieces of block per placement, usually spaced 15 to 30 feet apart. Maintain a constant supply of fresh bait for 10 days or until fresh signs of rat activity cease.

For burrow applications, insert one or two 2-oz. pieces of block deep into each active burrow. Check treated burrows each morning and evening to make sure that bait has not been pushed out of burrows.

For sewer applications, securely attach one end of a wire to bait block and the other end to a stationary structure such as the bottom step of manhole ladder or sewer grate, allowing just enough wire for bait to rest on a manhole bench. If no bench is present, suspend bait a few inches above the high water line, or place bait on a board supported by opposing steps of the ladder. Secure bait in this manner to minimize chance for removal by rats or water. Place at least one entire bait block per manhole.

Follow-up: Replace contaminated or spoiled bait immediately. Collect and dispose of all visible dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation occurs, repeat treatment. For a continuous infestation, set up permanent bait stations and replenish bait as needed. (15105)